PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5: (11) International Publication Number: WO 90/15593 A1 A61K 9/10, 9/14, 9/50 (43) International Publication Date: 27 December 1990 (27.12.90)

(21) International Application Number: PCT/SE90/00426 (74) Agent: AWAPATENT AB; Box 7045, S-103 94 Stockholm (SE).

(22) International Filing Date: 15 June 1990 (15.06.90) (81) Designated States: AT (European patent), AU, BE (Euro-

(30) Priority data: pean patent), BG, BR, CA, CH (European patent), DE (European patent)*, DK (European patent), ES (European patent), FI, FR (European patent), GB (European 8902257-8 21 June 1989 (21.06.89) SE patent), HU, IT (European patent), JP, KP, KR, LU (Eu-(71) Applicant (for all designated States except US): YTKEMISropean patent), NL (European patent), NO, RO, SE

KA INSTITUTET [SE/SE]; Box 5607, S-114 86 Stock-(European patent), SU, US. holm (SE).

(72) Inventors; and (75) Inventors; and
(75) Inventors/Applicants (for US only): SJÖSTRÖM, Brita [SE/SE]; Polhemsgatan 27¹¹, S-112 30 Stockholm (SE).
KRONBERG, Bengt [SE/SE]; Elsa Brändströms gata 19, S-126 63 Hägersten (SE). CARLFORS, Johan [SE/SE]; Prästgårdsvägen 13C, S-752 30 Uppsala (SE).
BLUTE, Irena [SE/SE]: Lojovägen 10, S-181 47 Liding (SE). Published

With international search report.

(54) Title: A PROCESS FOR THE PREPARATION OF DRUG PARTICLES

(57) Abstract

A process for the preparation of submicron size, monodisperse drug-particles of a drug of low water-solubility by emulsifying an organic solution of the drug in an aqueous phase and then removing the organic solvent resulting in drug precipitation, containing the steps: a) emulsifying the organic solution in the presence of an emulsifier comprising a surfactant capable of adsorption on the surface of a precipitated drug-particle: b) removing the organic solvent from the suspension; and c) recovering the precipitated drug-particles from the aqueous phase or storing the same in the original aqueous phase.

DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AU Australia FI Finland MG Madagascar BB Barbados FR France ML Mali BE Belgium GA Gabon MR Mauritania BF Burkina Fasso GB United Kingdom MW Malawi BG Bulgaria GR Greece NL Netherlands BJ Benin HU Hungary NO Norway BR Brazil IT Italy RO Romania CA Canada JP Japan SD Sudan CF Central African Republic KP Democratic People's Republic SE Sweden CG Congo Switzerland KR Republic of Korea SU Soviet Union CM Cameroon LI Liechtenstein TD Chad CG Germany, Federal Republic of LK Sri Lanka TG Iogo	AT Austria		ES	Spain	MC	Monaco
BB Barbados FR France ML Mali BE Belgium GA Gabon MR Mauritania BF Burkina Fasso GB United Kingdom MW Malawi BG Bulgaria GR Greece NL Netherlands BJ Benin HU Hungary NO Norway BR Brazil IT Italy RO Romania CA Canada JP Japan SD Sudan CF Central African Republic KP Democratic People's Republic SE Sweden CG Congo of Korea SN Sconegal CH Switzerland KR Republic of Korea SU Soviet Union CM Cameroon LI Liechtenstein TD Chad DE Germany, Federal Republic of LK Sri Lanka TG logo			Fi	Finland	MG	Madagascar
BE Belgium GA Gabon MR Mauritania BF Burkina Fasso GB United Kingdom MW Malawi BG Bulgaria GR Greece NL Netherlands BJ Benin HU Hungary NO Norway BR Brazil IT Italy RO Romania CA Canada JP Japan SD Sudan CF Central African Republic KP Democratic People's Republic SE Sweden CG Congo Switzerland KR Republic of Korea SU Soviet Union CM Cameroon LI Liechtenstein TD Chad CM Garmany, Federal Republic of LK Sri Lanka TG Fugo	_		FR	France	ML	
BF Burkina Fasso GB United Kingdom MW Malawi BG Bulgaria GR Greece NL Netherlands BJ Benin HU Hungary NO Norway BR Brazil IT Italy RO Romania CA Canada JP Japan SD Sudan CF Central African Republic KP Democratic People's Republic SE Sweden CG Congo KR Republic of Korea SN Scnegal CH Switzerland KR Republic of Korea SU Soviet Union CM Cameroon LI Licehtenstein TD Chad DE Germany, Federal Republic of LK Sri Lanka TG Fugo			GA	Gahon	MR	Mauritania
BG Bulgaria GR Greece NL Netherlands BJ Benin HU Hungary NO Norway BR Brazil IT Italy RO Romania CA Canada JP Japan SD Sudan CF Central African Republic KP Democratic People's Republic SE Sweden CG Congo of Korea SN Scnegal CH Switzerland KR Republic of Korea SU Soviet Union CM Cameroon L1 Liechtenstein TD Chad DE Germany, Federal Republic of LK Sri Lanka TG logo				United Kingdom	MW	Malawi
BJ Benin HU Hungary NO Norway BR Brazil IT Italy RO Romania CA Canada JP Japan SD Sudan CF Central African Republic KP Democratic People's Republic SE Sweden CG Congo of Korea SN Senegal CH Switzerland KR Republic of Korea SU Soviet Union CM Cameroon LI Liechtenstein TD Chad DE Germany, Federal Republic of LK Sri Lanka TG Fogo	-•			-	NL	Netherlands
BR Brazil IT Italy RO Romania CA Canada JP Japan SD Sudan CF Central African Republic KP Democratic People's Republic SE Sweden CG Congo of Korea SN Senegal CH Switzerland KR Republic of Korea SU Soviet Union CM Cameroon LI Liechtenstein TD Chad DE Germany, Federal Republic of LK Sri Lanka TG Fugo					NO	Norway
CA Canada JP Japan SD Sudan CF Central African Republic KP Democratic People's Republic SE Sweden CG Congo of Korea SN Senegal CH Switzerland KR Republic of Korea SU Soviet Union CM Cameroon LI Liechtenstein TD Chad DE Germany, Federal Republic of LK Sri Lanka TG Fugo	-				RO	Romania
CF Central African Republic KP Democratic People's Republic SE Sweden CG Congo of Korea SN Scinegal CH Switzerland KR Republic of Korea SU Soviet Union CM Cameroon L1 Liechtenstein TD Chad DE Germany, Federal Republic of LK Sri Lanka TG logo				_ =	SD	Sudan
CG Congo of Korea SN Senegal CH Switzerland KR Republic of Korea SU Soviet Union CM Cameroon LI Liechtenstein TD Chad DE Germany, Federal Republic of LK Sri Lanka TG logo		Aferma Dunublic	_	•	SE	Sweden
CH Switzerland KR Republic of Korea SU Soviet Union CM Cameroon LI Liechtenstein TD Chad DE Germany, Federal Republic of LK Sri Lanka TG logo			•••	• •	SN	Senegal
CM Cameroon LI Liechtenstein TD Chad DE Germany, Federal Republic of LK Sri Lanka TG logo	•		KR		SU	Soviet Union
DE Germany Federal Republic of LK Sri Lanka TG Fogo					TD	Chad
DE CHEMBEY, PEDERA REPORTE OF LACTOR LACTOR	_				TG	Fogo
DK Denmark LU Luxembourg US United States of Av	-	• .			us	United States of America

10

15

20

25

30

A process for the preparation of drug particles.

The present invention relates to a process for the preparation of submicron size, relatively monodisperse drug-particles of a drug of low water-solubility.

The size of water-insoluble drug-particles or such particles of a drug of low water-solubility is of considerable importance for the bioavailability of the drug. The rate of dissolution of the particles is directly proportional to their specific area and, accordingly, the size is critical for the absorption and distribution of the drug in the tissue. Another argument for producing submicron particles is that the smaller particle size will allow alternative ways of administration of the drug substance. For pharmaceutical applications drug-particles of submicron size are therefore desirable. Such particles are, in accordance with conventional techniques, produced by milling, a process which reduces the mean particle size to a few microns. However, the process produces a rather broad size-distribution thus resulting in illdefined dissolution kinetics. Such milling process frequently also results in contamination and degradation of the drug.

An alternative to such milling is to precipitate particles by crystallization. This is a more complex process which involves nucleation and growth kinetics but which is an alternative that overcomes the particle size limitations of milling processing. However, also according to this alternative the difficult inherent in using crystallization is that of controlling the particle size and the size distribution to meet with the requirements. Crystallization is a subject to several phenomena that aggravate the problem of controlling particle size. First, a crystal growth can uncontrollably change particle size and broaden the size distribution. Second, crystals formed may aggragate or agglomerate also resulting in uncontrolled size growth and distribution.

10

15

20

25

30

35

The present invention has for an object to provide a process for the preparation of small particles of water-in-soluble drugs or drugs of low water-solubility while controlling size distribution to avoid widely varying particle sizes.

Another object of the invention is to provide a process for such drug particle manufacture which is highly reproducible and easily controlled.

Yet another object of the invention is to device a process through which small drug particles can be produced which meet with the usual pharmaceutical requirements.

These and other objects of the invention will be clear from the following detail description. Accordingly, the invention resides in a process for the preparation of submicron size relatively monodisperse drug-particles of a drug of low water-solubility or a water-insoluble drug by emulsifying an organic solution of the drug in an aqueous phase and then removing the organic solvent resulting in drug precipitation. Such process comprises the following steps:

- a) emulsifying the organic solution in the presence of an emulsifier and if needed a protective colloid comprising a surfactant capable of adsorption on the surface of a precipitated drug particle;
- b) removing the organic solvent from the suspension; and
 c) recovering the precipitated drug particles from the aqueous phase or storing the same in the original aqueous phase.

Removal of the organic solvent from the suspension can take place in different ways. The following procedures are examples of such organic solvent removal:

the organic solvent can be removed by evaporation;
the organic solvent can be removed by washing through
continuous renewal of the aqueous phase;

the organic solvent can be removed by washing away same through dialysis by contact with a pure aqueous phase; the organic solvent can be removed by contacting the emulsion with an excess of aqueous phase to dissolve the or-

10

15

20

25

30

35

ganic solvent therein resulting in precipitation of the drug.

When removing the organic solvent by evaporation it may occur that water of the aqueous phase evaporates at the same time resulting in reduced water content of the emulsion. Such evaporation of water can be compensated for by adding further quantities of aqueous phase to the emulsion during evaporation of the organic solvent.

The different possibilities of controlling the drop diameter of the emulsion, and the drop size of the emulsion can be maintained within relatively broad range, such as between about 0.1 and about 20 mm. The size of the drops of the emulsion can be controlled by varying the supply of mechanic energy, by varying the concentration of surfactant and by varying the ratio between organic solvent and aqueous phase in the emulsion. The concentration of surfactant can for practical reasons be molarly related to the molar concentration of the drug.

The surfactant used is subject to several requirements as to its function. Thus, it shall operate satisfactorily as an emulsifying agent, i.e. it shall result in the formation of a stable emulsion. It is possible to use a protective colloid in combination with the surfactant to improve the stability of the emulsion. The basic requirement according to the invention is to use a surfactant which is capable of adsorption on the surface of a precipitated drug-particle. By using such surfactants a controlled and reproducible precipitation of small drug-particles takes place and the particle size distribution will be maintained within fairly narrow ranges.

Furthermore, the surfactant preferably inhibits aggregation and agglomeration of the particles and should advantageously be capable of inhibiting crystal growth. Such crystal growth is partly eliminated in view of reduced surface energy when the surfactant is adsorbed on the surface of the crystals.

It is preferred to use a protective colloid to stabilize the suspension. Said colloid can be synthetic, semisynthetic or can be constituted by a polysaccharide or a pro-

10

15

tein. Parts of the protective colloid must be water-soluble to give colloidal stability. Micell formation in solution should be avoided but this is not necessary if the solubilizing capacity thereof on the crystal-forming substance is low. Possible incorporation of the tenside and/or the protective colloid in the crystal structure of the particle is permissible.

It is important to note that the process of the present invention does not involve any reaction in the system and that the process is not based on so called micro-emulsions.

It is advantageous in the process of the invention that no or very little mass transportation takes place between the drops of the emulsion and this requires a certain minimum concentration of drug in said drops.

Below there are given non-limiting examples of drugs, organic solvents and surfactants suitable for use in the process of the invention.

20 DRUGS

As water-insoluble drugs or drugs of low water-solubility the following are of interest.

Examples of drugs whose bioavailability has been increased as a result of particle size reduction

25	a result of particle size reduction				
	A, vitamin	Medroxyprogesterone acetate			
	4-Acetamidophenyl,2,2,2-tri-	Nitrofurantoin			
	-chlorethyl carbonate	Phenobarbital			
	Aspirin	Phenacetin			
30	Bishydroxycoumarin	Procaine penicillin			
	Chloramphenicol	Reserpine			
	Cyheptamide	Spironolactone			
	Digoxin	Sulfadiazine			
	Fluocinolone acetonide	Sulfasoxazole			
35	Griseofulvin	Sulfur			
	p-Hydroxypropiophenone	Tolbutamide			

	Drugs with potential bioequivalency problems.				
	Acetazolamide	Para-aminosalicylic acid			
	Acetyldigitoxin	Para-methadione			
	Alseroxylon	Perphenazine			
5	Aminophyllin	Phenacemide			
_	Aminosalicylic acid	Phensuximide			
	Bendroflumethiazide	Phenylaminosalicylate			
	Benzthiazide	Phenytoin			
	Betamethasone	Phytonadione			
10	Bishydroxycoumarin	Polythiazide			
	Chlorambucil	Prednisolone			
	Chlorodiazepoxide	Primidone			
	Chlorothiazide	Probenecid			
	Chloropromazine	Procainamide			
15	Cortisone acetate	Prochlorperazine			
	Deserpidine	Promazine			
	Dexamethasone	Promethazine			
	Dichlorphenamide	Propylthiouracil			
•	Dienestrol	Pyrimethamine			
20	Diethylstilbestrol	Quinethiazide			
	Dyphylline	Quinidine			
	Ethinyl estradiol	Rauwolfia serpentina			
	Ethosuxmide	Rescinnamine			
	Ethotoin	Reserpine			
25	Ethoxzolamide	Salicylazosulfapyridine			
	Fludrocortisone	Sodium sulfoxone			
	Fluphenazine	Spironolactone			
	Fluprednisolone	Sulfadiazine			
	Hydralazine	Sulfadimethoxine			
30	Hydrochlorothiazide	Sulfamerazine			
	Hydroflumethiazide	Sulfaphenazole			
-	Imipramine	Sulfasomidine			
	Isoproterenol	Sulfasoxazole			
	Liothyronine	Theophylline			
35	Menadione	Thioridazine			
	Mephenytoin	Tolbutamide			

cont.

Methazolamide Triamcinolone

Methyclothiazide Trichlormethiazide

Methylprednisolone Triethyl melamine

Methyltestosterone Trifluoperazine

Nitrofurantoin Triflupromazine

Oxtriphylline Trimeprazine

Trimethadione

Uracil mustand

10 .

5

ORGANIC SOLVENTS

Among suitable organic solvents for use in the emulsion the following can be mentioned.

Warfarin

Any organic solvent that is a liquid and is poorly soluble in water. A prerequisite is that the solvent must be removed to concentrations that are acceptable from a toxicological point of view.

20 Examples:

- linear, branched or cyclic alkanes with carbon number of 5 mor higher; e.g. n-pentane, n-heptane and higher linear alkanes; 2,2-dimethyl butane or 2,2,4-trimethyl pentane; cyclopentane, cyclohexane, methylcyclohexane etc.
- 25 linear, branched or cyclic alkenes with carbon number of 5 or higher; e.g. 2-pentene, cyclohexene, 1,3 cyclopentadiene or 2,3 dimethyl-1-pentene.
 - linear, branched or cyclic alkynes with carbon number of 5 or higher; e.g. 2-pentyne or 4-methyl-2-pentyne.
- aromatic hydrocarbons; e.g. toluene, ethylbenzene.

 completely or partially halogenated hydrocarbons; e.g.

 dichlormetan, chloroform, chlorobenzene, chlorobenzoic acid

 etc.
 - ethers; e.g. diethylether etc.
- esters; e.g. ethylacetate, 9-Octadecenoic acid, ethyl ester, ethyl oleate, tetradecanoic acid 1-methylethyl ester, etc.

- ketones; e.g. cyclohexanone, 2-pentanone etc.
- mono-, di- or tri-glycerides; e.g. synthetic glycerol triacetate, glycerolmonolinolein etc., and native oils: almond oil, cotton seed oil, corn oil, olive oil, peanut oil, sesame oil, soybean oil etc.
- alcohols; e.g. benzenemethanol pentanol, hexanol etc.
- aldehydes; e.g. hexanol,
- acids; e.g. hexanoic acid,
- amines; e.g. 1-aminoheptane,
- 10 nitriles; e.g. amylcyanide,
 - silicones, linear or cyclic; e.g. octamethyltetrasiloxane or hexamethyldisiloxane,
 - or any combination of these derivatives; e.g. 1-chloro-3-ethylhexane,
- 15 or any combination of these solvents

SURFACTANTS

Surfactants or groups of surfactants meeting with the requirements as indicated above are for example:

20

25

30

5

Glyceryl monoalkylate; diacetyl tartaric acid esters of monoand diglycerides of edible fats or oils, or edible fat-forming fatty acids; mono- and diglycerides of edible fats or oils, or edible fatforming acids; monosodium phosphate derivatives of mono- and diglycerides of edible fats or oils, or edible fat-forming fatty acids; glycerol ester of wood rosin; alcyl monoglyceridyl citrate;

succistearin (stearoyl propylene glycol hydrogen succinate); dioctyl sodium sulfosuccinate; lecithins (pure and technical qualities); hydroxylated lecithins; methyl glucoside-coconut oil ester; sodium alcyl sulfate; potassium alcyl sulfate; sodium or potassium mono- and dimethyl naphthalene sulfonates; sodium or potassium alcyl fumarate; acetylated monogly-

10

cerides; succinylated monoglycerides; monoglyceride citrate; ethoxylated mono- and diglycerides; ethoxylated sorbitan esters; sorbitan monostearate; sorbitan esters of fatty acids; calcium alcyloyl-2-lactylate; sodium alcyloyl-2-lactylate; potassium alcyloyl-2-lactylate; lactylic esters of fatty acids; lactylated fatty acid esters of glycerol and propylene glycol; glyceryl-lacto esters of fatty acids; ethoxylated alcyl phenols and alcohols; polyglycerol esters of fatty acids; propylene glycol mono- and diesters of fats and fatty acids; sucrose fatty acid esters; fatty acids; salts of fatty acids; synthetic fatty alcohols; poloxamers; meroxapol; poloxamine; "pluradot", or any combination of these.

PROTECTIVE COLLOIDS

In the following there are also given examples of preferred protective colloids:

Gums: e.g. acacia, agar, carrageenan, guar, karaya, locust bean, pectin, propylene glycol alginate, sodium alginate, tragacanth, xanthan, gum arabicum.

- Cellulosics: e.g. carboxymethylcellulose, sodium, microcrystalline cellulose and carboxymethylcellulose, sodium, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropyl methylcellulose, cellulose acetate phtalate, methylcellulose, ethyl hydroxyethyl cellulose, methocel.
- Clays: e.g. Bentonite (colloidal aluminum silicate), colloidal magnesium aluminum silicate (hectorite), colloidal magnesium aluminum silicate (attapulgite), magnesium silicate (sepiolite).
- Miscellaneous: e.g. Carbomer NF, gelatin, polyethylene glycols, polypropyleleglycols and copolymers thereof, lecithins,
 Carbopol 934, Veegum, polyacrylic acid, polymethacrylic acid,
 polyacrylic acid-CO-acrylamide, polyvinylpyrrolidon, polyvinylalcohol with varying degree of hydrolysis.

Proteins: e.g. albumine, gelatine, casein or any combination
of these

The invention will be further described in the following by non-limiting examples. In these examples the drug model used is cholesteryl acetate, which substance is useful for illustrating the problem solved by the present invention.

In these examples reference is made to the appended drawings, wherein:

Fig. 1 illustrates the effect of the method of emulsification on the particle size;

Fig. 2 illustrates a diagram on the size of emulsion droplets and the cholesterol acetate particles as a function of the concentration of cholesterol acetate in toluene; and

Fig. 3 illustrates the size of the emulsion droplets and the cholesterol acetate particles as a function of the concentration of surfactant.

15

20

25

30

35

10

5

EXAMPLE 1

An emulsion of cholesteryl acetate dissolved in toluene and an aqueous phase containing ethoxylated nonylphenol ether as a surfactant is prepared in the following manner. The drug model substance, cholesteryl acetate, is dissolved in toluene. The solution is emulsified with an aqueous phase containing ethoxylated nonylphenol ether as a surfactant to form an oil-in-water-type emulsion.

The organic solvent, toluene, is then evaporated from the emulsion, whereby the drug model substance precipitates and the crystals are stabilized by the surfactant, said surfactant being adsorbed on the surface of the precipitated particles.

Using the emulsion ingredients indicated above a series of tests are made to establish the effect of the method of emulsification on the particle size, the influence of the concentration of the drug substance in toluene on the size of the particles and the influence of concentration of surfactant in the emulsion on the size of the emulsion droplets and thus also on the size of the model drug-particles.

15

20

25

30

This illustration of different effects is made in relation to the appended drawings numbered 1 to 3 and related to the respective effects outlined above. In Fig. 1 the particles were obtained from emulsions which were prepared by vibration, and the particle sizes were measured by an electrozone sensing method. In Fig. 2 the emulsions were prepared by homogenization with a microfluidizer, and the particle sizes were measured by quasi-elastic light scattering. With regard to Fig. 3 the emulsions were prepared by homogenization with a microfluidizer.

Yet another alternative to control the emulsion droplet size is to vary the oil/water phase ratio.

EXAMPLE 2

This example illustrates the use of a surfactant capable of stabilizing the emulsion and suspension of cholesterol acetate and cyclohexane, respectively. The emulsion is homogenized in a microfluidizer. The oil/water phase ratio is 10/90, and the amount of surfactant is 5% by weight based on the weight of the oil phase. The surfactant used is a mixture of Tween 80 and Span 80 at a weight ratio of 1:9. Span 80 is a tradename for sorbitan monooleate and Tween 80 a tradename of POE-(20)-sorbitan monooleate.

The particle size in the suspension lies within the range of between about 80 nm and about 400 nm. The particle size is measured after 10 days and is found to be about constant.

EXAMPLE 3

Example 2 is repeated but using a mixture of Tween 20 and Span 80 (weight ratio 16:1) as a surfactant. Similar results are obtained.

EXAMPLE 4

Example 2 is repeated but using Tween 80 as a surfactant.

EXAMPLE 5

Example 2 is repeated but using DK ESTER F-160 (a sucrose ester) as a surfactant.

5

EXAMPLE 6

Example 2 is repeated but using Ovothin 170 (egg phospholipid) as a surfactant.

10 EXAMPLE 7

Example 2 is repeated but using Epikuran 145 (soybean lecithin) as a surfactant.

EXAMPLE 8

Example 2 is repeated but using Epikuran 200 SH (soybean lecithin) and the sodium salt of glycocholic acid (4:1) as a surfactant.

EXAMPLE 9

20 Example 2 is repeated but using Epikuran 200 SH and the sodium salt of taurocholic acid (4:1) as a surfactant.

EXAMPLE 10

Example 2 is repeated but using Ovothin 170 and the sodium salt of glycocholic acid (4:1) as a surfactant.

EXAMPLE 11

Example 2 is repeated but using Ovothin 170 and the sodium salt of taurocholic acid (4:1) as a surfactant.

30

EXAMPLE 12

Example 2 is repeated but using Triodan 55, polyglycerol esters of fatty acids as a surfactant.

35 EXAMPLE 13

Example 2 is repeated but using Acidan N-12, citric acid esters of monoglycerides as a surfactant.

EXAMPLE 14

Example 2 is repeated but using Epikuran 200 SH as a surfactant and polyvinylpyrrolidone as a protective colloid.

In the above Examples 4-14 results corresponding to those of Example 2 are obtained.

10

15

20

25


30


35


CLAIMS

- 1. A process for the preparation of submicron size, relatively monodisperse drug-particles of a drug of low water-solubility by emulsifying an organic solution of the drug in an aqueous phase and then removing the organic solvent resulting in drug precipitation, characterized by the steps:
- a) emulsifying the organic solution in the presence of an emulsifier, and if needed a protective colloid, comprising a surfactant capable of adsorption on the surface of a precipitated drug-particle;
- b) removing the organic solvent from the suspension; and
 c) recovering the precipitated drug-particles from the
 aqueous phase or storing the same in the original aqueous
 phase.
- 2. A process according to claim 1, characterized in that the organic solvent is removed by evaporation.
- 3. A process according to claim 1, characterized in that the organic solvent is removed by washing through continuous renewal of the aqueous phase.
- 4. A process according to claim 1, characterized in that the organic solvent is removed by washing away same through dialysis by contact with a pure aqueous phase.
- 5. A process according to claim 1, characterized in that the organic solvent is removed by contacting the emulsion with an excess of aqueous phase to dissolve the organic solvent therein resulting in precipitation of the drug.
- 6. A process according to claim 2, wherein aqueous phase is added during evaporation to compensate for co-evaporation of water from the aqueous phase.
- 7. A process according to any preceding claim, wherein the drop diameter of the emulsion is controlled by the concentration of surfactant.
- 8. A process according to claim 7, wherein said concentration is related to the molar concentration of the drug.
- 9. A process according to any preceding claim, wherein the surfactant is pharmaceutically acceptable.

10. Drug particles whenever prepared by the process of any preceding claim.







*Calculated with respect to CA only.

INTERNATIONAL SEARCH REPORT

International Application No PCT/SE 90/00426

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶						
According to International Patent Classification (IPC) or to both National Classification and IPC						
IPC5: I	IPC5: A 61 K 9/10, 9/14, 9/50					
II. FIELD	II. FIELDS SEARCHED					
		Minimum Docume				
Classificat	on Sys	em C	Classification Symbols			
	·					
IPC5		A 61 K; A 61 J; B 01 J				
		Documentation Searched other to the Extent that such Document	r than Minimum Documentation s are included in Fields Searched ⁸			
		,				
SE,DK,	FI,NO) classes as above				
III. DOCU	MENTS	CONSIDERED TO BE RELEVANT 9				
Category *	C	itation of Document, ¹¹ with indication, where ap	propriate, of the relevant passages 12	Relevant to Claim No. ¹³		
Х	US,	A, 4826689 (M.R. VIOLANTO E	T AL.)	1,3,4,5,		
		2 May 1989,	4 3ima 2.	9,10		
		see column 2, line 60 - column 5, lines 7-36	umn 4, 11ne 2;			
		Corumn 5, Times 7 50				
X	EP,	A, 0169618 (STERILIZATION T	ECHNICAL SERVICES	1,3-5,9-		
		INC) 29 January 1986,	5 line 10	10		
	}	see page 4, line 22 - page	s, The 19			
A	US,	A, 3979520 (W. ROTHE ET AL.)	1,2,6,		
		7 September 1976,		10		
		see the whole document				
Α	DE,	A, 2817453 (ROHNER AG PRATT	ELN)	1-10		
	}	31 October 1979,	0 lima 14			
		see page 7, line 7 - page 1	o, line 14			
						
* Special categories of cited documents: 10 "T" later document published after the international filing date or priority date and not in conflict with the application but						
considered to be of particular relevance invention						
"E" earlier document but published on or after the international filing date "X" document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve a tree inventive step.						
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the						
"O" document referring to an oral disclosure, use, exhibition or						
other means in the art.						
"P" document published prior to the international filing date but later than the priority date claimed "&" document member of the same patent family IV. CERTIFICATION						
Date of the Actual Completion of the International Search Date of Mailing of this International Search Report						
12th September 1990						
International Searching Authority Signature of Authorized Officer						
SWEDISH PATENT OFFICE NIKTAS FORSTUNG						

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.PCT/SE 90/00426

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the Swedish Patent Office EDP file on 90-08-02 The Swedish Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report		Publication date	Patent family member(s)		Publication date	
US-A-	4826689	89-05-02	NONE			
EP-A-	0169618	86-01-29	AU-B- AU-D- PT-A-	579166 4501285 80494	88-11-17 87-01-22 85-11-22	
US-A-	3979520	76-09-07	AT-B- AU-D- BE-A- CA-A- DE-A-C- FR-A-B- GB-A- JP-C- JP-B- JP-B- NL-A- SE-B-C- SE-A- US-A-	330954 7365374 820254 1029661 2348334 2244455 1420907 1089246 50058216 56032967 7412393 402210 7412044 4060634	76-07-26 76-04-01 75-03-24 78-04-18 75-03-27 75-04-18 76-01-14 82-03-23 75-05-21 81-07-31 75-04-01 78-06-26 75-03-27 77-11-29	
DE-A-	2817453	79-10 - 31	NONE		_	